

# **ECHO LAKE 2018**

## **Lake Stewardship Monitoring Report**

King County Water & Land Resources Division  
Science & Technical Support Section  
[www.kingcounty.gov/EnvironmentalScience](http://www.kingcounty.gov/EnvironmentalScience)

---

### **Summary & Recommendations**

*Thank you to Saul Malamud and the City of Shoreline staff for monitoring Echo Lake.*

#### **The key takeaways from the 2018 monitoring season are:**

- Nitrogen, phosphorus, and chlorophyll concentrations continued to decline from peaks in 2015.
- Even with the recent declines, Echo Lake's nutrient concentrations and algal growth were still fairly high, and the water was less clear.
- Echo Lake's high nutrient concentrations and nitrogen-to-phosphorus (N:P) ratios below 25 both indicate that Echo Lake is likely to have algal blooms dominated by cyanobacteria (which have the ability to produce toxins).
- An algal bloom was sampled for toxin testing in late October. Toxin testing found microcystin present at 5.0 µg/L, below the Washington State Recreational Guideline of 6.0 µg/L.

#### **The Lake Stewardship Program recommends:**

- Explore what has helped to decrease nutrient concentrations in Echo Lake since 2015 – and encourage those trends to continue. Reducing nutrient concentrations will likely help to reduce algal blooms.
  - Stay alert for toxic algae blooms in Echo Lake – increase people's awareness of toxic algae, and their ability to identify which algae are potentially toxic. Any potentially toxic blooms should be reported to the King County Lake Stewardship Program and sampled for toxin analysis.
  - Monitoring is a key part of good lake stewardship, building a valuable long-term dataset to guide lake management and detect any future problems. Continue to monitor Echo Lake through the Lake Stewardship Program.
-

### ***In this report:***

- **What We Measure & Why**
- **Water Quality Results & Trends**
  - 2018 monitoring results
  - Long-term annual averages
  - Trends over time
- **Trophic State**
  - Trophic state indices
  - Comparison map
- **Supplemental Data**
  - Summary statistics
  - Hydrology: Lake level and precipitation
  - Year-round Secchi depth and water temperature
  - Water column profile
  - Total alkalinity
  - Water color

## **What We Measure & Why**

- **Secchi depth** is a measure of water clarity or transparency. Secchi depth is shallower when there are more suspended particles in the lake, such as sediment or algae. Secchi depth is also affected by water color, often from tannins or other naturally occurring organic molecules.
- **Water temperature** can affect the growth rates of plants and algae. In addition, cooler or warmer water temperatures favor different species of fish and other aquatic organisms.
- **Chlorophyll-a** is a measure of the amount of algae in a lake. Chlorophyll-a is a pigment necessary for algae to photosynthesize and store energy.
- **Phosphorus** and **nitrogen** are naturally occurring nutrients necessary for growth and reproduction in both plants and animals. Increases in nutrients (especially phosphorus) can lead to more frequent and dense algal blooms.
- The **ratio of total nitrogen to total phosphorus (N:P)** indicates whether nutrient conditions favor the growth of cyanobacteria (blue-green algae). When N:P ratios are near or below 25, cyanobacteria can dominate the algal community. This is important because cyanobacteria have the ability to produce toxins.

## **Water Quality Results & Trends**


The following graphs show the water-quality parameters that are sampled from May through October, at 1 m depth (additional depths and parameters are measured on profile days; see *Supplemental Data*). The left column of graphs shows results for each sampling

date in 2018, and the right column shows average values for each year (May-October averages).


Data for Echo Lake are the blue circles (with white centers) connected by the blue line. Any gaps in the blue line indicate missed samples. To provide some context for these values, the grey points in the background are results for all other lakes in the Lake Stewardship program.

Any long-term trends in Echo Lake are drawn with a dashed red line and described further after the graphs. Statistical trend analyses used a seasonal (monthly) Kendall test ( $p < 0.05$ ).

### 2018 Monitoring Results



### Long-Term Annual Averages



Echo Lake's high nutrient concentrations and nitrogen-to-phosphorus (N:P) ratios below 25 both indicate that Echo Lake is likely to have algal blooms dominated by cyanobacteria (which have the ability to produce toxins).

The table below gives more details about the long-term trends. Results are presented as an average amount and percent of change per decade (the increase or decrease over ten years). Percent change is calculated as the percent of the estimated value in 2001, when monitoring started.

Parameter	Change per Decade	(%)
Total phosphorus	6.5 µg/L	(23%)
N:P ratio	-4.6	(-20%)


## Trophic State

The Trophic State Index (TSI) is a common index of a lake’s overall biological productivity. TSI values are calculated from Secchi depth, chlorophyll-a concentrations, and total phosphorus concentrations. These three TSI estimates are all scaled between 0 and 100.

TSI calculations use average values from June-September, focusing on fairly consistent “summer” conditions. This is in contrast with the annual averages shown above, which also include May and October data.

*Oligotrophic* lakes (TSI <40) are very clear, with low nutrient concentrations and low algal growth. *Eutrophic* lakes (TSI >50) have less-clear water, with high nutrient concentrations and high algal growth. Eutrophic lakes are more likely to have frequent algal blooms. *Mesotrophic* lakes (TSI 40-50) are in the middle, with fairly clear water, and moderate nutrient concentrations and algal growth. Lakes in lowland King County have a range of different natural trophic states, and human activities may also alter a lake’s trophic state (usually by changing nutrient inputs).


## Trophic state indices



In 2018, the TSI values were in the eutrophic or upper mesotrophic range.

## Comparison map

For a comparison with other lakes, this map shows the trophic state for each lake in the King County Lake Stewardship program in 2018. The color of each circle indicates the lake's average chlorophyll-a TSI value for the year.



## Supplemental Data


### Summary statistics

This table summarizes data from 2018 (1 m depth only), giving the minimum, mean (average), and maximum values for each parameter. This includes summary statistics for the full 2018 calendar year for Secchi and temperature, which were measured year-round, and May-October summary statistics for all parameters. To reduce biases from missing data or changes in sampling frequency, monthly means were calculated and then averaged to give an overall mean.

Parameter	Minimum	Mean	Maximum
<b>Full-year statistics</b>			
Secchi depth (m)	0.3	2.3	3.9
Water temperature (°C)	5.0	13.3	24.0
<b>May-October statistics</b>			
Secchi depth (m)	0.3	2.3	3.9
Water temperature (°C)	12.0	18.8	24.0
Chlorophyll-a (µg/L)	2.0	10.7	23.2
Total nitrogen (µg/L)	360.0	495.4	850.0
Total phosphorus (µg/L)	19.9	31.7	47.3
N:P ratio	9.2	16.5	27.9


### Hydrology: Lake level and precipitation

Lake level and precipitation were recorded year-round. Bars show total weekly precipitation, and the line shows average weekly lake level.



## Year-round Secchi depth and water temperature

Secchi depth and water temperature (at 1 m depth) were measured weekly in 2018. The blue circles (with white centers) and blue line are data for Echo Lake. Gaps in the line indicate missed sampling dates. Grey points in the background are results for all other lakes in the Lake Stewardship program.



Note that Echo Lake's year-round measurements were taken from a dock where the water was about 1.5 m deep. Only Secchi depths shallower than 1.5 m could be measured from this location.

## Water column profile

In May and August, water was collected at the mid-lake sampling station from three depths in a water-column profile: 1 m, the middle depth of the water column, and 1 m from the lake bottom.


Date	Depth	Temp	Chlor	Pheo	TN	NH3	NO2/3	TP	OPO4
5/21/2018	1.0	19.2	2.7	(1.3)	360	33.2	12.0	39.1	7.4
	3.5	14.7	1.3	(1.4)	496	–	–	62.4	–
	7.0	7.9	1.9	5.5	922	394.0	236.0	171.0	105.0
8/20/2018	1.0	22.1	6.1	1.3	434	2.4	(10.0)	19.9	1.2
	3.5	22.0	7.4	1.6	436	–	–	20.2	–
	7.0	9.3	150.0	–	1470	540.0	(10.0)	336.0	37.1

Parameter abbreviations are: chlorophyll-a (Chlor), pheophytin (Pheo), total nitrogen (TN), ammonia (NH3), nitrate/nitrite (NO2/3), total phosphorus (TP), orthophosphate (OPO4). Depth is in m, temperature is in °C, and all other parameters are in µg/L. Dashes indicate parameters that were not analyzed for a given sample. Values below the method detection limit (MDL) are enclosed in parentheses and have the value of the MDL substituted.

## Total alkalinity

A lake’s ability to resist acidification, also called its buffering capacity, is measured as “total alkalinity.” Lakes with total alkalinity less than 20 mg CaCO<sub>3</sub> are considered sensitive to acidification. We measured total alkalinity in May and August (on profile-sampling days) at 1 m depth. In 2018, the average total alkalinity of these two samples was 16.4 mg CaCO<sub>3</sub>.

The blue circles (with white centers) and blue line are annual average alkalinity values for Echo Lake. Grey points in the background are results for all other lakes in the Lake Stewardship program.




## Water color

Water color affects a lake’s water clarity (and Secchi depth). Water color is measured by shining a specific wavelength of ultraviolet light (254 nm) through a filtered water sample

and measuring the percent that was absorbed. We measured UV254 absorbance in May and August (on profile-sampling days) at 1 m depth. In 2018, the average UV254 absorbance of these two samples was 0.09, on a scale where 0 is no absorbance (perfectly clear) and 1 is complete absorbance (perfectly opaque).

The blue circles (with white centers) and blue line are annual average UV absorbance values for Echo Lake. Grey points in the background are results for all other lakes in the Lake Stewardship program.



**King County**

Department of Natural Resources and Parks  
Water and Land Resources Division  
**Science and Technical Support Section**  
King Street Center, KSC-NR-0600  
201 South Jackson Street, Suite 600  
Seattle, WA 98104  
206-477-4800 TTY Relay: 711  
[www.kingcounty.gov/EnvironmentalScience](http://www.kingcounty.gov/EnvironmentalScience)