

# **LAKE FENWICK 2021**

## **Lake Stewardship Monitoring Report**

King County Water & Land Resources Division  
Science & Technical Support Section  
[www.kingcounty.gov/EnvironmentalScience](http://www.kingcounty.gov/EnvironmentalScience)

---

### **Summary & Recommendations**

*Thank you to TJ Neff, the volunteer monitor for Lake Fenwick.*

#### **The key takeaways from the 2021 monitoring season are:**

- Lake Fenwick has fairly clear water with high nutrient concentrations and high algal growth.
- In comparison to other lakes in the program, Lake Fenwick experienced some of the highest nutrient concentrations in 2021.
- Nitrogen-to-phosphorus (N:P) ratios were below 25 for much of the monitoring season. This indicates the potential for algal blooms to be dominated by cyanobacteria (which have the ability to produce toxins).
- Total phosphorus exceeded the lake's TMDL (total maximum daily load) of 19 µg/L during each sampling event.
- An algal bloom was sampled for toxin testing in September. No algal toxins were detected in this sample.
- Lake Fenwick was last monitored in the Lake Stewardship Program from 2001 - 2004.

#### **The Lake Stewardship Program recommends:**

- Stay alert for toxic algae blooms in Lake Fenwick – increase people's awareness of toxic algae, and their ability to identify which algae are potentially toxic. Any potentially toxic blooms should be reported to the King County Lake Stewardship Program and sampled for toxin analysis.
  - Monitoring is a key part of good lake stewardship, building a valuable long-term dataset to guide lake management and detect future problems. Continue to monitor Lake Fenwick through the Lake Stewardship Program.
-

## **What We Measure & Why**

- **Secchi depth** is a measure of water clarity or transparency. Secchi depth is shallower when there are more suspended particles in the lake, such as sediment or algae. Secchi depth is also affected by water color, often from tannins or other naturally occurring organic molecules.
- **Water temperature** can affect the growth rates of plants and algae. In addition, cooler or warmer water temperatures favor different species of fish and other aquatic organisms. Many lakes in King County naturally have cold water, so increases in water temperature can favor non-native species.
- **Chlorophyll-a** is a measure of the amount of algae in a lake. Chlorophyll-a is a pigment necessary for algae to photosynthesize and store energy.
- **Phosphorus** and **nitrogen** are naturally occurring nutrients necessary for growth and reproduction in both plants and animals. Increases in nutrients (especially phosphorus) can lead to more frequent and dense algal blooms.
- The **ratio of total nitrogen to total phosphorus (N:P)** indicates whether nutrient conditions favor the growth of cyanobacteria (blue-green algae). When N:P ratios are near or below 25, cyanobacteria can dominate the algal community. Tracking this ratio is important because cyanobacteria have the ability to produce toxins.


## **Water Quality Results & Trends**

The following graphs show the water-quality parameters that are sampled from May through October, at 1 m depth (additional depths and parameters are measured on profile days; see *Supplemental Data*). The left column of graphs shows results for each sampling date in 2021, and the right column shows average values (May-October) for each year the lake was monitored in the Lake Stewardship Program.


Results for Lake Fenwick are the blue circles connected by the blue line. Gaps in the blue line indicate missed samples. The grey points in the background are results for all other lakes in the [Lake Stewardship program](#). These points provide some context for Lake Fenwick values.

Long-term trends in Lake Fenwick, if present, are shown with a dashed red line and described below the graphs. Statistical trend analyses use a seasonal (monthly) Kendall test ( $p < 0.05$ ).

## 2021 Monitoring Results



## Long-Term Annual Averages



Nitrogen-to-phosphorus (N:P) ratios were below 25 for much of the monitoring season. This indicates the potential for algal blooms to be dominated by cyanobacteria (which have the ability to produce toxins).

The table below gives more details about the long-term trends. Results are presented as an average amount and percent of change per decade (the increase or decrease over ten years). Percent change is calculated as the percent of the estimated value in 2001, when monitoring started.

Parameter	Change per Decade	(%)
Total nitrogen	64 µg/L	(13%)

Total nitrogen concentrations have increased over time. However, with the gap in monitoring years and few years of monitoring this trend is less reliable.

## Trophic State


The Trophic State Index (TSI) is a common index of a lake’s overall biological productivity. Index values are calculated from Secchi depth, chlorophyll-a concentrations, and total phosphorus concentrations, all scaled between 0 and 100.

TSI calculations use average values from June-September, focusing on fairly consistent “summer” conditions. This is in contrast with the annual averages shown above, which also include May and October data. Based on TSI values, a lake is usually classified as being in one of the three possible classes: oligotrophic, mesotrophic, or eutrophic.

- *Oligotrophic* lakes (TSI <40) are very clear, with low nutrient concentrations and low algal growth.
- *Mesotrophic* lakes (TSI 40-50) have fairly clear water, and moderate nutrient concentrations and algal growth.
- *Eutrophic* lakes (TSI >50) have less-clear water, with high nutrient concentrations and high algal growth. Eutrophic lakes are also more likely to have frequent algal blooms.

Lakes in lowland King County have a range of different natural trophic states, and human activities may also alter a lake’s trophic state (usually by changing nutrient inputs).


## **Trophic state indices**



In 2021, the chlorophyll and total phosphorus TSI values were in the eutrophic range while the Secchi TSI value was in the mesotrophic range.

## Comparison map

For a comparison with other lakes, the following map shows the trophic state for each lake in the King County Lake Stewardship program in 2021. The color of each circle indicates the lake's average chlorophyll-a TSI value for the year.



## Supplemental Data

### Summary statistics

The following table summarizes data from May-October 2021 (1 m depth only), giving the minimum, mean (average), and maximum values for each parameter. To reduce biases from missing data or changes in sampling frequency, monthly means were calculated and then averaged to give an overall mean.

Parameter	Minimum	Mean	Maximum
Secchi depth (m)	3.5	4.3	5.7
Water temperature (°C)	15.0	19.6	22.7
Chlorophyll-a (µg/L)	1.7	16.1	72.3
Total nitrogen (µg/L)	445.0	632.5	928.0
Total phosphorus (µg/L)	26.1	39.0	74.3
N:P ratio	8.7	17.9	31.1

### Water column profile


In May and August, water was collected at the mid-lake sampling station from three depths in a water column profile: 1 m, the middle depth of the water column, and 1 m from the lake bottom.

Date	Depth	Temp	Chlor	Pheo	TN	NH3	NO2/3	TP	OPO4
5/16/2021	1	17.0	8.9	1.1	580	6.7	(10.0)	51.9	2.0
	4	13.5	11.8	1.7	510	–	–	36.5	–
	8	10.0	12.2	1.5	522	92.7	42.4	88.1	9.1
8/8/2021	1	22.6	1.7	1.4	445	10.5	(10.0)	26.1	0.6
	4	16.2	2.7	1.7	621	–	–	48.5	–
	8	8.5	56.9	–	1410	927.0	(10.0)	528.0	361.0

Parameter abbreviations are: chlorophyll-a (Chlor), pheophytin (Pheo), total nitrogen (TN), ammonia (NH3), nitrate/nitrite (NO2/3), total phosphorus (TP), orthophosphate (OPO4). Depth is in m, temperature is in °C, and all other parameters are in µg/L. Dashes indicate parameters that were not analyzed for a given sample. Values below the method detection limit (MDL) are enclosed in parentheses and have the value of the MDL substituted.

## Water column temperatures


Temperature was measured at all depths for every event in 2021. A lake experiences thermal stratification when surface and bottom depth temperatures are different. This is typically seen in the summer when surface temperatures are warmer than bottom temperatures. In the fall as the weather conditions cool down, surface temperatures decrease and the lake mixes as the bottom water and surface water are closer in temperatures.



In 2021, Lake Fenwick was thermally stratified during the summer, while mixing is seen in the beginning and end of the monitoring season.

## Dissolved oxygen

Dissolved oxygen (DO) was collected at profile depths at every event in Lake Fenwick using a YSI Professional Optical Dissolved Oxygen instrument. During summer thermal stratification, concentrations of DO at the bottom of the lake become anoxic, meaning devoid of oxygen. This occurs when algae and other organic matter settles to the bottom of the lake and decompose, which consumes oxygen. DO concentrations are higher in the surface and the middle of the lake due to DO produced by photosynthesizing algae.




In 2021, Lake Fenwick remained anoxic at its middle and bottom depths for most of the year.

## Total alkalinity

A lake’s ability to resist acidification, also called its buffering capacity, is measured as “total alkalinity.” Lakes with total alkalinity less than 20 mg CaCO<sub>3</sub> are considered sensitive to acidification. We measured total alkalinity in May and August (on profile-sampling days) at 1 m depth. In 2021, the average total alkalinity of these two samples was 38.9 mg CaCO<sub>3</sub>.


The blue circles and blue line are annual average alkalinity values for Lake Fenwick. Grey points in the background are results for all other lakes in the Lake Stewardship program.



## Water color


Water color affects a lake's water clarity (and Secchi depth). Water color is measured by shining a specific wavelength of ultraviolet light (254 nm) through a filtered water sample and measuring the percent that was absorbed. We measured UV254 absorbance in May and August (on profile sampling days) at 1 m depth. In 2021, the average UV254 absorbance of these two samples was 0.15, on a scale where 0 is no absorbance (perfectly clear) and 1 is complete absorbance (perfectly opaque).

The blue circles (with white centers) and blue line are annual average UV absorbance values for Lake Fenwick. Grey points in the background are results for all other lakes in the Lake Stewardship program.



## Lake inlet

Water from the Lake Fenwick inlet was sampled and analyzed for total phosphorus to assess the impact of external loading on lake water quality. Samples were collected once a month.



Visit the [King County Lake Stewardship](#) website for more data and information.



**King County**

Department of Natural Resources and Parks  
Water and Land Resources Division  
Science and Technical Support Section  
King Street Center, KSC-NR-5600  
201 South Jackson Street, Suite 5600  
Seattle, WA 98104  
206-477-4800 TTY Relay: 711  
[www.kingcounty.gov/EnvironmentalScience](http://www.kingcounty.gov/EnvironmentalScience)